

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
26 February 2004 (26.02.2004)

PCT

(10) International Publication Number  
**WO 2004/016088 A2**

(51) International Patent Classification<sup>7</sup>: **A01N 43/40**,  
C07D 213/61

(21) International Application Number:  
PCT/EP2003/009516

(22) International Filing Date: 8 August 2003 (08.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
02356159.0 12 August 2002 (12.08.2002) EP  
0305233 29 April 2003 (29.04.2003) FR

(71) Applicant (*for all designated States except US*): **BAYER CROPSCIENCE S.A.** [FR/FR]; 16, rue Jean-Marie Leclair, F-69009 LYON (FR).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MANSFIELD, Darren, James** [GB/FR]; 25, montée Bonafous, F-69004 Lyon (FR). **COOKE, Tracey** [GB/GB]; 7 Larch Avenue, Brickett Wood, AL2, St Albans AL2 3SN (GB). **THOMAS, Peter, Stanley** [GB/GB]; 94 Balsham Road, Linton, Cambridge, Cambridgeshire CB1 6LW (GB). **COQUERON, Pierre-Yves** [FR/FR]; 36, rue de la Tête d'Or, F-69006 Lyon (FR). **VORS, Jean-Pierre** [FR/FR]; 16, chemin de Montpellas, F-69009 Lyon (FR). **BRIGGS, Geoffrey, Gower** [GB/GB]; 46 Granby Avenue, Harpenden, Herts, Hertfordshire AL5 5QR (GB). **LACHAISE, Hélène** [FR/FR]; 11, chemin de Montpellas, F-69009 Lyon (FR). **RIECK, Heiko** [DE/FR]; 9, rue Claude Monet, F-69110 Sainte Foy Les Lyon (FR). **DESBORDES, Philippe**

[FR/FR]; 30, rue Bancel, F-69007 Lyon (FR). **GROS-JEAN-COURNOYER, Marie-Claire** [FR/FR]; La Trève Collonge, Route d'Albigny, F-69250 Curis au Mont d'Or (FR).

(74) Agent: **MERIGEAULT, Shona**; Bayer Cropscience S.A., 14-20 rue Pierre Baizet, BP 9163, F-69263 LYON CEDEX 09 (FR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**


— *of inventorship (Rule 4.17(iv)) for US only*

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: NOVEL 2-PYRIDYLETHYLBENZAMIDE DERIVATIVE



(57) Abstract: Compound of general formula (I) in which p is an integer equal to 1, 2, 3 or 4; q is an integer equal to 1, 2, 3, 4 or 5; each substituent X is chosen, independently of the others, as being halogen, alkyl or haloalkyl, at least one of the substituents being a haloalkyl; each substituent Y is chosen, independently of the others, as being halogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl,

halothioalkyl, alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or benzylsulphonyl; as to the N-oxides of 2-pyridine thereof; with the exception of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-ethyl}-2,6-dichlorobenzamide. Method for preparing the compound of general formula (I). Fungicidal composition comprising the compound of general formula (I). Method for treating phytopathogenic diseases.

**Novel 2-pyridylethylbenzamide derivative**


The present invention relates to novel fungicidal compounds, to the method  
5 for preparing them, to the fungicidal compositions comprising these compounds, and  
also to their use in the agricultural field as fungicides.

Patent Application WO 01/11965 describes a broad family of fungicidal  
compounds of general formula encompassing the compounds of the present  
invention. However, the said compounds are not described in that patent application  
10 and their activity as a fungicide was not tested.

It is however always useful, in the agricultural field, to use compounds which  
are more active than those already known to those skilled in the art, with the aim of  
decreasing the amounts of active material to be used by the farmer, whilst at the  
same time maintaining an effectiveness at least equivalent to the already known  
15 compounds.

It has now been discovered that a certain number of compounds, selected  
from a broad family of compounds, possess the above mentioned advantages.

A subject of the present invention is therefore a family of fungicidal  
20 compounds of general formula (I):



in which :

- p is an integer equal to 1, 2, 3 or 4;
  - q is an integer equal to 1, 2, 3, 4 or 5;
  - each substituent X is chosen, independently of the others, as being halogen, alkyl or  
25 haloalkyl, at least one of the substituents being a haloalkyl;
  - each substituent Y is chosen, independently of the others, as being halogen, alkyl,  
alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl,  
cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl,  
alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or  
30 benzylsulphonyl;
- as to the N-oxides of 2-pyridine thereof;

with the exception of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2,6-dichlorobenzamide.

In the context of the present invention, the substituents X of the 2-pyridine and the substituents Y of the benzene ring will be indexed in order to facilitate understanding. Thus, for example, if p is equal to 2 and q is equal to 1, the substituents termed "X" will be denoted by X<sup>1</sup> and X<sup>2</sup> and the substituent termed "Y" will be denoted by Y<sup>1</sup>.


For the purposes of the present invention, halogen means chlorine, bromine, iodine or fluorine.

For the purposes of the present invention, each of the alkyl or acyl radicals present in the molecule contains from 1 to 10 carbon atoms, preferentially from 1 to 7 carbon atoms, more preferentially from 1 to 5 carbon atoms, and may be linear or branched.

For the purposes of the present invention, each of the alkenyl or alkynyl radicals present in the molecule contains from 2 to 10 carbon atoms, preferentially from 2 to 7 carbon atoms, more preferentially from 2 to 5 carbon atoms, and may be linear or branched.


The present invention relates to compound of general formula (I). Preferentially, compounds of general formula (I) have the following characteristics, taken individually or in combination:

- p is chosen equal to 2, the substituents X<sup>1</sup> and X<sup>2</sup> being positioned as follows:




- q is chosen equal to 1 or 2, the substituent(s) Y being positioned in the ortho position of the benzene ring.

A preferred subfamily of compounds according to the invention consists of the compounds corresponding to general formula (I')



X and Y being as defined above. More preferably,  $X^1$  is chosen as being halogen and  $X^2$  is chosen as being haloalkyl.

Another preferred subfamily of compounds according to the invention consists of the compounds corresponding to general formula (I''):



5 the substituents X and Y being as defined above. More preferably, compound of general formula (I'') according to the present invention have the following characteristics, taken individually or in combination:

- $X^1$  is chosen as being halogen and  $X^2$  is chosen as being haloalkyl;
- $Y^1$  is chosen as being halogen or haloalkyl.

10 More preferably, the haloalkyl group is chosen as being trifluoromethyl.


Even more preferably, a subject of the present invention is the following compounds:

- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide;
- 15 - N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide;
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-bromobenzamide.

20 The present invention also relates to a process for the preparation of the compound of general formula (I). Thus, according to a further aspect of the present invention there is provided a process for the preparation of compound of formula (I) as defined above which comprises:

- a first step consisting in reacting, in the presence of a base in aprotic polar solvent, a compound of general formula (Ia) in order to substitute it selectively in the 2-position:

\* either with a group of the alkyl cyanoacetate type (NC-CH<sub>2</sub>-CO<sub>2</sub>Alk) to produce a compound of general formula (Ib) according to the following reaction scheme:



where: - X is as previously defined;

5


- Alk represents an alkyl radical;

- Q is a nucleofugal radical, preferably chosen to be a halogen

or trifluoromethanesulphonate;

the compound of general formula (Ib) thus obtained then undergoing dealkyloxy-carbonylation in the presence of an alkali metal halide, such as Li-halogen, K-halogen or Na-halogen, at the reflux of a water/dimethyl sulphoxide mixture, according to the Krapcho reaction described in A.P. *Synthesis*, **1982**, 805, 893, to produce the compound of general formula (Ic) according to the following reaction scheme:


10



sodium halide will preferentially be used in the context of the present invention;

15


\* or with acetonitrile to directly produce the compound of general formula (Ic) according to the following reaction scheme:




- a second step consisting in the reduction of the compound of general formula (Ic) to pyridylethanamine of general formula (Id) (or its corresponding ammonium salt depending on whether or not the medium is acid) under hydrogen pressure in the presence of a metal catalyst in a protic solvent according to the following reaction scheme:

20





- a third step consisting in converting the compound of general formula (Id) to a compound of general formula (I) by reaction with a benzoyl halide in the presence of a base according to the following reaction scheme:



The second step of the above described process is conducted in the presence of a metal catalyst. Preferably, the metal catalyst is chosen as being a nickel-, platinum- or palladium-based catalyst.

The third step of the above mentioned process is conducted in the presence of a benzoyl halide. Preferably, the benzoyl halide is chosen as being benzoyl chloride.

From compound of general formula (I) obtained by virtue of the method of preparation described above, those skilled in the art will be able to prepare, by virtue of methods known to them, the N-oxide derivatives of 2-pyridine. For example, the compound of general formula (I) obtained by virtue of the method according to the present invention may be treated with an excess of meta-chloroperbenzoic acid (also called m-CPBA) in the presence of a solvent, which may be chloroform, at a temperature which may range from 60 to 80°C.

The present invention also relates to a fungicidal composition comprising an effective amount of an active material of general formula (I). Thus, according to the present invention, there is provided a fungicidal composition comprising, as an active ingredient, an effective amount of a compound of general formula (I) as defined above and an agriculturally acceptable support, carrier or filler.

In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid.

5 Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

The composition may also comprise additional components. In particular, the composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential

10 when the active material and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content may be comprised between 5% and 40% by weight of the composition.

Optionally, additional components may also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active materials can be combined with any

25 solid or liquid additive, which complies with the usual formulation techniques.

In general, the composition according to the invention may contain from 0.05 to 99% (by weight) of active material, preferably 10 to 70% by weight.

Compositions according to the present invention can be used in various forms

30 such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable

35 concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment, seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed

treatment, suspension concentrate (flowable concentrate), ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

5        These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which must be diluted before application to the crop.

10       The compounds of the invention can also be mixed with one or more insecticides, fungicides, bactericides, attractant acaricides or pheromones or other compounds with biological activity. The mixtures thus obtained have a broadened spectrum of activity. The mixtures with other fungicides are particularly advantageous.

15       The fungicidal compositions of the present invention can be used to curatively or preventively control the phytopathogenic fungi of crops. Thus, according to a further aspect of the present invention, there is provided a method for curatively or preventively controlling the phytopathogenic fungi of crops characterised in that a fungicidal composition as hereinbefore defined is applied to the seed, the plant and/or to the fruit  
20 of the plant or to the soil in which the plant is growing or in which it is desired to grow.

The composition as used against phytopathogenic fungi of crops comprises an effective and non-phytotoxic amount of an active material of general formula (I).

25       The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the fungi present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be controlled, the type of crop, the climatic conditions and the compounds included in the fungicidal composition according to the invention.

30       This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

35       The method of treatment according to the present invention is useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.



Among the plants that can be protected by the method according to the invention, mention may be made of cotton; flax; vine; fruit crops such as *Rosaceae* sp. (for instance pip fruits such as apples and pears, but also stone fruits such as apricots, almonds and peaches), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp.,  
 5 *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantains), *Rubiaceae* sp., *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruits); leguminous crops such as *Solanaceae* sp. (for instance tomatoes), *Liliaceae* sp., *Asteraceae* sp. (for instance lettuces), *Umbelliferae* sp., *Cruciferae* sp.,  
 10 *Chenopodiaceae* sp., *Cucurbitaceae* sp., *Papilionaceae* sp. (for instance peas), *Rosaceae* sp. (for instance strawberries); big crops such as *Graminae* sp. (for instance maize, cereals such as wheat, rice, barley and triticale), *Asteraceae* sp. (for instance sunflower), *Cruciferae* sp. (for instance colza), *Papilionaceae* sp. (for instance soja), *Solanaceae* sp. (for instance potatoes), *Chenopodiaceae* sp. (for  
 15 instance beetroots); horticultural and forest crops; as well as genetically modified homologues of these crops.

Among the plants and the possible diseases of these plants protected by the method according to the present invention, mention may be made of :

- wheat, as regards controlling the following seed diseases: fusaria  
 20 (*Microdochium nivale* and *Fusarium roseum*), stinking smut (*Tilletia caries*, *Tilletia controversa* or *Tilletia indica*), septoria disease (*Septoria nodorum*) and loose smut;
- wheat, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yallundae*, *Tapesia acuiformis*), take-all (*Gaeumannomyces graminis*), foot blight (*F. culmorum*, *F. graminearum*), black speck  
 25 (*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts (*Puccinia striiformis* and *Puccinia recondita*) and septoria diseases (*Septoria tritici* and *Septoria nodorum*);
- wheat and barley, as regards controlling bacterial and viral diseases, for example barley yellow mosaic;
- 30 - barley, as regards controlling the following seed diseases: net blotch (*Pyrenophora graminea*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut (*Ustilago nuda*) and fusaria (*Microdochium nivale* and *Fusarium roseum*);
- barley, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yallundae*), net blotch (*Pyrenophora teres* and  
 35 *Cochliobolus sativus*), powdery mildew (*Erysiphe graminis forma specie hordei*), dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);

- potato, as regards controlling tuber diseases (in particular *Helminthosporium solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*), mildew (*Phytophthora infestans*) and certain viruses (virus Y);

5 - potato, as regards controlling the following foliage diseases: early blight (*Alternaria solani*), mildew (*Phytophthora infestans*);

- cotton, as regards controlling the following diseases of young plants grown from seeds: damping-off and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*) and black root rot (*Thielaviopsis basicola*);

10 - protein yielding crops, for example peas, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*, *Mycosphaerella pinodes*), fusaria (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*) and mildew (*Peronospora pisi*);

- oil-bearing crops, for example rape, as regards controlling the following seed diseases: *Phoma lingam*, *Alternaria brassicae* and *Sclerotinia sclerotiorum*;

15 - corn, as regards controlling seed diseases: (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp., and *Gibberella fujikuroi*);

- flax, as regards controlling the seed disease: *Alternaria linicola*;

- forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*);

20 - rice, as regards controlling the following diseases of the aerial parts: blast disease (*Magnaporthe grisea*), bordered sheath spot (*Rhizoctonia solani*);

- leguminous crops, as regards controlling the following diseases of seeds or of young plants grown from seeds: damping-off and collar rot (*Fusarium oxysporum*, *Fusarium roseum*, *Rhizoctonia solani*, *Pythium* sp.);

25 - leguminous crops, as regards controlling the following diseases of the aerial parts: grey mould (*Botrytis* sp.), powdery mildews (in particular *Erysiphe cichoracearum*, *Sphaerotheca fuliginea* and *Leveillula taurica*), fusaria (*Fusarium oxysporum*, *Fusarium roseum*), leaf spot (*Cladosporium* sp.), alternaria leaf spot (*Alternaria* sp.), anthracnose (*Colletotrichum* sp.), septoria leaf spot (*Septoria* sp.),  
30 black speck (*Rhizoctonia solani*), mildews (for example *Bremia lactucae*, *Peronospora* sp., *Pseudoperonospora* sp., *Phytophthora* sp.);

- fruit trees, as regards diseases of the aerial parts: monilia disease (*Monilia fructigenae*, *M. laxa*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*);

- vine, as regards diseases of the foliage: in particular grey mould (*Botrytis cinerea*), powdery mildew (*Uncinula necator*), black rot (*Guignardia biwelli*) and mildew (*Plasmopara viticola*);

5 - beetroot, as regards the following diseases of the aerial parts: cercospora blight (*Cercospora beticola*), powdery mildew (*Erysiphe beticola*), leaf spot (*Ramularia beticola*).

The fungicide composition according to the present invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber"  
10 means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists in contacting one or more compounds of the present invention, or a composition according to the invention; this includes for example direct application, spraying,  
15 dipping, injection or any other suitable means.

The dose of active material usually applied in the treatment according to the present invention is generally and advantageously between 10 and 800 g/ha, preferably between 50 and 300 g/ha for applications in foliar treatment. The dose of active  
20 substance applied is generally and advantageously between 2 and 200 g per 100 kg of seed, preferably between 3 and 150 g per 100 kg of seed in the case of seed treatment. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to adapt the application doses according to the nature of the crop to be treated.

25

The fungicidal composition according to the present invention may also be used in the treatment of genetically modified organisms with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into whose genome a heterologous  
30 gene encoding a protein of interest has been stably integrated. The expression "heterologous gene encoding a protein of interest" essentially means genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.

35


The compositions according to the present invention may also be used for the preparation of composition useful to curatively or preventively treat human and animal

fungus diseases such as, for example, mycoses, dermatoses, trichophyton diseases and candidiasis or diseases caused by *Aspergillus spp.*, for example *Aspergillus fumigatus*.

5           The aspects of the present invention will now be illustrated with reference to the following tables of compounds and examples. The following Tables A and B illustrate in a non-limiting manner examples of fungicidal compounds according to the present invention. In the following Examples, M+1 means the molecular ion peak, plus or minus 1 a.m.u. (atomic mass units) respectively, as observed in mass  
10 spectroscopy and M (ApCI+) means the molecular ion peak as it was found via positive atmospheric pressure chemical ionisation in mass spectroscopy.



Table A



Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-1	H	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	H	H	363
A-2	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	H	363
A-3	Cl	CF <sub>3</sub>	H	H	H	H	H	Cl	Cl	397
A-4	Cl	CF <sub>3</sub>	H	H	F	H	H	H	H	347
A-5	Cl	CF <sub>3</sub>	H	H	F	H	H	F	H	365
A-6	Cl	CF <sub>3</sub>	H	H	Cl	H	H	Cl	H	397
A-7	Cl	CF <sub>3</sub>	H	H	OMe	H	H	H	H	359
A-8	Cl	CF <sub>3</sub>	H	H	OMe	OMe	H	H	H	389
A-9	Cl	CF <sub>3</sub>	H	H	Me	H	H	H	H	343
A-10	Cl	CF <sub>3</sub>	H	H	H	H	F	H	H	347
A-11	Cl	CF <sub>3</sub>	H	H	H	H	Cl	H	H	363
A-12	Cl	CF <sub>3</sub>	H	H	H	H	OMe	H	H	359
A-13	Cl	CF <sub>3</sub>	H	H	H	H	OMe	OMe	H	389
A-14	Cl	CF <sub>3</sub>	H	H	H	H	OMe	H	OMe	389

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-15	Cl	CF <sub>3</sub>	H	H	H	H	H	F	H	347
A-16	Cl	CF <sub>3</sub>	H	H	H	H	H	OMe	H	359
A-17	Cl	CF <sub>3</sub>	H	H	H	H	H	OBu	H	401
A-18	Cl	CF <sub>3</sub>	H	H	H	H	H	C(CH <sub>3</sub> ) <sub>3</sub>	H	397
A-19	Cl	CF <sub>3</sub>	H	H	H	H	H	Me	H	343
A-20	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	H	H	397
A-21	Cl	CF <sub>3</sub>	H	H	H	H	H	Et	H	357
A-22	Cl	CF <sub>3</sub>	H	H	H	H	H	Pr	H	371
A-23	Cl	CF <sub>3</sub>	H	H	H	H	H	Bu	H	385
A-24	Cl	CF <sub>3</sub>	H	H	H	H	H	C <sub>5</sub> H <sub>11</sub>	H	399
A-25	Cl	CF <sub>3</sub>	H	H	F	F	H	H	H	365
A-26	Cl	CF <sub>3</sub>	H	H	H	H	Me	H	H	343
A-27	Cl	CF <sub>3</sub>	H	H	H	H	CH <sub>2</sub> Cl	H	H	377
A-28	Cl	CF <sub>3</sub>	H	H	H	H	CF <sub>3</sub>	H	H	397
A-29	Cl	CF <sub>3</sub>	H	H	NO <sub>2</sub>	H	H	H	H	374
A-30	Cl	CF <sub>3</sub>	H	H	H	H	NO <sub>2</sub>	H	H	374
A-31	Cl	CF <sub>3</sub>	H	H	H	H	H	NO <sub>2</sub>	H	374
A-32	Cl	CF <sub>3</sub>	H	H	H	H	Cl	H	Cl	396
A-33	Cl	CF <sub>3</sub>	H	H	F	H	H	H	F	365
A-34	Cl	CF <sub>3</sub>	H	H	H	H	H	F	F	365
A-35	Cl	CF <sub>3</sub>	H	H	H	H	F	H	F	365

14

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-36	Cl	CF <sub>3</sub>	H	H	H	H	CN	H	H	354
A-37	Cl	CF <sub>3</sub>	H	H	Me	Me	H	Me	H	371
A-38	Cl	CF <sub>3</sub>	H	H	H	H	NO <sub>2</sub>	Me	H	388
A-39	Cl	CF <sub>3</sub>	H	H	H	H	H	CO <sub>2</sub> Me	H	387
A-40	Cl	CF <sub>3</sub>	H	H	F	F	F	H	H	383
A-41	Cl	CF <sub>3</sub>	H	H	F	H	H	F	F	383
A-42	Cl	CF <sub>3</sub>	H	H	F	F	H	F	H	383
A-43	Cl	CF <sub>3</sub>	H	H	F	H	F	F	F	401
A-44	Cl	CF <sub>3</sub>	H	H	F	Cl	H	H	H	382
A-45	Cl	CF <sub>3</sub>	H	H	I	H	H	H	H	455
A-46	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	H	407
A-47	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	F	H	H	413
A-48	Cl	CF <sub>3</sub>	H	H	Cl	H	Cl	H	H	397
A-49	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	Cl	397
A-50	Cl	CF <sub>3</sub>	H	H	NHCH <sub>3</sub>	H	H	H	H	358
A-51	Cl	CF <sub>3</sub>	H	H	OMe	H	OMe	H	H	389
A-52	Cl	CF <sub>3</sub>	H	H	OH	H	H	H	H	345
A-53	Cl	CF <sub>3</sub>	H	H	OH	H	Me	H	H	359
A-54	Cl	CF <sub>3</sub>	H	H	OH	H	H	Cl	H	379
A-55	Cl	CF <sub>3</sub>	H	H	OH	H	H	OMe	H	375
A-56	Cl	CF <sub>3</sub>	H	H	OH	H	H	Me	H	359

15

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-57	Cl	CF <sub>3</sub>	H	H	OH	H	H	H	F	363
A-58	Cl	CF <sub>3</sub>	H	H	OH	H	H	H	OMe	375
A-59	Cl	CF <sub>3</sub>	H	H	OH	H	H	H	Me	359
A-60	Cl	CF <sub>3</sub>	H	H	Me	H	Me	H	H	357
A-61	Cl	CF <sub>3</sub>	H	H	Me	H	H	Me	H	357
A-62	Cl	CF <sub>3</sub>	H	H	Me	H	H	H	Me	357
A-63	Cl	CF <sub>3</sub>	H	H	OH	H	Cl	H	H	379
A-64	Cl	CF <sub>3</sub>	H	H	F	H	Cl	H	H	381
A-65	Cl	CF <sub>3</sub>	H	H	OH	F	H	H	H	363
A-66	Cl	CF <sub>3</sub>	H	H	Me	H	Cl	H	H	377
A-67	Cl	CF <sub>3</sub>	H	H	Me	H	F	H	H	361
A-68	Cl	CF <sub>3</sub>	H	H	OH	H	H	F	H	363
A-69	Cl	CF <sub>3</sub>	H	H	F	H	H	H	Cl	381
A-70	Cl	CF <sub>3</sub>	H	H	Me	H	OH	H	H	359
A-71	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	OMe	436
A-72	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	H	CF <sub>3</sub>	465
A-73	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	F	H	415
A-74	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	465
A-75	Cl	CF <sub>3</sub>	H	H	F	CF <sub>3</sub>	Cl	H	H	449
A-76	Cl	CF <sub>3</sub>	H	H	PhCH <sub>2</sub>	H	H	H	H	419
A-77	Cl	CF <sub>3</sub>	H	H	Me	H	NO <sub>2</sub>	H	H	388



16

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-78	Cl	CF <sub>3</sub>	H	H	Me	H	NO <sub>2</sub>	H	NO <sub>2</sub>	433
A-79	Cl	CF <sub>3</sub>	H	H	Me	H	H	H	NO <sub>2</sub>	388
A-80	Cl	CF <sub>3</sub>	H	H	Me	H	H	Br	H	421
A-81	Cl	CF <sub>3</sub>	H	H	Me	H	OMe	H	H	373
A-82	Cl	CF <sub>3</sub>	H	H	OMe	H	OMe	OMe	H	419
A-83	Cl	CF <sub>3</sub>	H	H	OMe	H	H	OMe	H	389
A-84	Cl	CF <sub>3</sub>	H	H	OMe	H	H	OMe	OMe	419
A-85	Cl	CF <sub>3</sub>	H	H	OMe	H	H	H	OMe	389
A-86	Cl	CF <sub>3</sub>	H	H	OE <sub>t</sub>	H	H	H	H	373
A-87	Cl	CF <sub>3</sub>	H	H	OMe	H	H	Cl	H	393
A-88	Cl	CF <sub>3</sub>	H	H	OMe	H	H	H	Cl	393
A-89	Cl	CF <sub>3</sub>	H	H	OCF <sub>3</sub>	H	H	H	H	413
A-90	Cl	CF <sub>3</sub>	H	H	OMe	H	H	H	F	377
A-91	Cl	CF <sub>3</sub>	H	H	OPr	H	H	H	H	387
A-92	Cl	CF <sub>3</sub>	H	H	OMe	H	H	H	t-Bu	415
A-93	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	Br	441
A-94	Cl	CF <sub>3</sub>	H	H	I	H	I	H	I	397
A-95	Cl	CF <sub>3</sub>	H	H	Cl	H	NO <sub>2</sub>	H	H	408
A-96	Cl	CF <sub>3</sub>	H	H	Cl	H	H	NO <sub>2</sub>	H	408
A-97	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	NO <sub>2</sub>	408
A-98	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	SO <sub>2</sub> F	445

12

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-99	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	SMe	409
A-100	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	Cl	441
A-101	Cl	CF <sub>3</sub>	H	H	Cl	H	NO <sub>2</sub>	H	Cl	442
A-102	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	Br	485
A-103	Cl	CF <sub>3</sub>	H	H	Cl	H	Cl	H	Cl	431
A-104	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	CF <sub>3</sub>	431
A-105	Cl	CF <sub>3</sub>	H	H	Br	H	NO <sub>2</sub>	H	H	452
A-106	Cl	CF <sub>3</sub>	H	H	Cl	H	H	Cl	F	415
A-107	Cl	CF <sub>3</sub>	H	H	Cl	H	H	F	F	399
A-108	Cl	CF <sub>3</sub>	H	H	Cl	H	OMe	OMe	H	423
A-109	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	NO <sub>2</sub>	452
A-110	Cl	CF <sub>3</sub>	H	H	Br	H	H	Cl	H	441
A-111	Cl	CF <sub>3</sub>	H	H	Br	H	Cl	H	H	441
A-112	Cl	CF <sub>3</sub>	H	H	F	Br	H	H	H	425
A-113	Cl	CF <sub>3</sub>	H	H	Cl	F	F	H	H	399
A-114	Cl	CF <sub>3</sub>	H	H	Et	H	H	H	H	357
A-115	Cl	CF <sub>3</sub>	H	H	Me	H	H	Me	CO <sub>2</sub> Et	429
A-116	Cl	CF <sub>3</sub>	H	H	Me	H	Me	H	Cl	391
A-117	Cl	CF <sub>3</sub>	H	H	Me	H	H	F	H	361
A-118	Cl	CF <sub>3</sub>	H	H	Me	H	Cl	F	H	395

18

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-119	Cl	CF <sub>3</sub>	H	H	O-CF <sub>2</sub> - CHFCI	H	H	Cl	H	495
A-120	Cl	CF <sub>3</sub>	H	H	O-allyl	H	H	Cl	H	419
A-121	Cl	CF <sub>3</sub>	H	H	O-allyl	H	H	H	H	385
A-122	Cl	CF <sub>3</sub>	H	H	Br	H	F	H	H	425
A-123	Cl	CF <sub>3</sub>	H	H	Cl	H	Me	F	Me	409
A-124	Cl	CF <sub>3</sub>	H	H	Cl	F	Me	H	H	395
A-125	Cl	CF <sub>3</sub>	H	H	Cl	H	F	Cl	H	415
A-126	Cl	CF <sub>3</sub>	H	H	Cl	H	H	CF <sub>3</sub>	H	431
A-127	Cl	CF <sub>3</sub>	H	H	SMe	H	H	Cl	H	409
A-128	Cl	CF <sub>3</sub>	H	H	SO <sub>2</sub> Me	H	H	Cl	H	441
A-129	Cl	CF <sub>3</sub>	H	H	Br	H	H	NO <sub>2</sub>	H	452
A-130	Cl	CF <sub>3</sub>	H	H	Cl	H	H	SO <sub>2</sub> Me	H	441
A-131	Cl	CF <sub>3</sub>	H	H	Cl	H	H	CN	H	388
A-132	Cl	CF <sub>3</sub>	H	H	Cl	H	Cl	Cl	H	431
A-133	Cl	CF <sub>3</sub>	H	H	SO <sub>2</sub> Me	H	H	Br	H	485
A-134	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	Me	421
A-135	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	SO <sub>2</sub> NMe <sub>2</sub>	H	504
A-136	Cl	CF <sub>3</sub>	H	H	Cl	H	OH	SO <sub>2</sub> Me	H	457
A-137	Cl	CF <sub>3</sub>	H	H	Br	H	H	SO <sub>2</sub> Me	H	485

19

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-138	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	SO <sub>2</sub> Me	H	475
A-139	Cl	CF <sub>3</sub>	H	H	SO <sub>2</sub> Me	H	H	CF <sub>3</sub>	H	475
A-140	Cl	CF <sub>3</sub>	H	H	SMe	H	H	F	H	393
A-141	Cl	CF <sub>3</sub>	H	H	H	H	Br	Cl	H	441
A-142	Cl	CF <sub>3</sub>	H	H	Me	H	H	SMe	H	389
A-143	Cl	CF <sub>3</sub>	H	H	SMe	H	H	CF <sub>3</sub>	H	443
A-144	Cl	CF <sub>3</sub>	H	H	SMe	H	Cl	Cl	H	443
A-145	Cl	CF <sub>3</sub>	H	H	Cl	H	H	SMe	H	409
A-146	Cl	CF <sub>3</sub>	H	H	SMe	H	Br	Cl	H	487
A-147	Cl	CF <sub>3</sub>	H	H	H	H	CF <sub>3</sub>	Cl	H	431
A-148	Cl	CF <sub>3</sub>	H	H	SMe	H	OMe	F	H	423
A-149	Cl	CF <sub>3</sub>	H	H	F	H	OMe	F	H	395
A-150	Cl	CF <sub>3</sub>	H	H	Br	H	H	SMe	H	453
A-151	Cl	CF <sub>3</sub>	H	H	iPr	H	Br	H	H	449
A-152	Cl	CF <sub>3</sub>	H	H	Et	H	H	Br	H	435
A-153	Cl	CF <sub>3</sub>	H	H	SMe	H	H	Br	H	453
A-154	Cl	CF <sub>3</sub>	H	H	Br	H	OMe	Br	H	515
A-155	Cl	CF <sub>3</sub>	H	H	SO <sub>2</sub> Me	H	F	Cl	H	459
A-156	Cl	CF <sub>3</sub>	H	H	NO <sub>2</sub>	H	H	Me	Me	402
A-157	Cl	CF <sub>3</sub>	H	H	SMe	H	O-CH <sub>2</sub> - CF <sub>2</sub>	Br	H	533




Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-158	Cl	CF <sub>3</sub>	H	H	H	H	Br	I	H	533
A-159	Cl	CF <sub>3</sub>	H	H	F	H	SO <sub>2</sub> NMe <sub>2</sub>	F	H	472
A-160	Cl	CF <sub>3</sub>	H	H	H	H	NO <sub>2</sub>	Me	Me	402
A-161	Cl	CF <sub>3</sub>	H	H	F	H	H	SO <sub>2</sub> Me	H	425
A-162	Cl	CF <sub>3</sub>	H	H	Me	H	F	Cl	H	395
A-163	Cl	CF <sub>3</sub>	H	H	Br	H	F	CF <sub>3</sub>	H	493
A-164	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	OCF <sub>3</sub>	491
A-165	Cl	CF <sub>3</sub>	H	H	F	H	SMe	F	H	411
A-166	Cl	CF <sub>3</sub>	H	H	CHCH <sub>2</sub>	H	H	F	H	373
A-167	Cl	CF <sub>3</sub>	H	H	SMe	H	H	H	H	375
A-168	Cl	CF <sub>3</sub>	H	H	F	H	SMe	Br	H	471
A-169	Cl	CF <sub>3</sub>	H	H	H	H	I	OH	I	597
A-170	Cl	CF <sub>3</sub>	H	H	CN	H	H	H	H	354
A-171	Cl	CF <sub>3</sub>	H	H	OMe	H	H	NO <sub>2</sub>	H	404
A-172	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	H	F	415
A-173	Cl	CF <sub>3</sub>	H	H	F	CF <sub>3</sub>	H	H	H	415
A-174	Cl	CF <sub>3</sub>	H	H	F	I	H	H	H	473
A-175	Cl	CF <sub>3</sub>	H	H	Br	H	H	H	F	425
A-176	Cl	CF <sub>3</sub>	H	H	I	H	H	Cl	H	489
A-177	Cl	CF <sub>3</sub>	H	H	I	H	H	H	Me	469
A-178	Cl	CF <sub>3</sub>	H	H	Cl	H	Me	H	H	377

21

Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
A-179	Cl	CF <sub>3</sub>	H	H	I	H	Me	H	H	469
A-180	Cl	CF <sub>3</sub>	H	H	Br	H	Me	H	H	421
A-181	Cl	CF <sub>3</sub>	H	H	Cl	H	H	OMe	OMe	423
A-182	Cl	CF <sub>3</sub>	H	H	F	Cl	Me	H	H	395
A-183	Cl	CF <sub>3</sub>	H	H	I	H	H	H	Br	533
A-184	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	Me	377
A-185	Cl	CF <sub>3</sub>	H	H	Cl	H	H	H	I	489
A-186	Cl	CF <sub>3</sub>	H	H	Me	H	H	H	F	361
A-187	Cl	CF <sub>3</sub>	H	H	OCHF <sub>2</sub>	H	H	H	H	395
A-188	Cl	CF <sub>3</sub>	H	H	I	H	H	H	Cl	489
A-189	Cl	CF <sub>3</sub>	H	H	I	H	H	OMe	OMe	515
A-190	Cl	CF <sub>3</sub>	H	H	Br	H	H	F	H	425
A-191	Cl	CF <sub>3</sub>	H	H	CHF <sub>2</sub>	H	H	H	H	422
A-192	Cl	CF <sub>3</sub>	H	H	S-CHF <sub>2</sub>	H	H	H	H	411
A-193	Cl	CF <sub>3</sub>	H	H	Me	H	NH <sub>2</sub>	H	H	358
A-194	Cl	CF <sub>3</sub>	H	H	NH <sub>2</sub>	H	Me	H	H	358
A-195	Cl	CF <sub>3</sub>	H	H	iPr	H	H	H	H	371
A-196	Cl	CF <sub>3</sub>	H	H	H	H	H	CF <sub>3</sub>	H	397


Table B



Composé n°	X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Y <sup>1</sup>	Y <sup>2</sup>	Y <sup>3</sup>	Y <sup>4</sup>	Y <sup>5</sup>	M+1
B-1	Cl	CF <sub>3</sub>	H	H	CF <sub>3</sub>	H	H	H	H	413
B-2	Cl	CF <sub>3</sub>	H	H	OMe	H	H	H	H	375
B-3	Cl	CF <sub>3</sub>	H	H	F	F	H	HI	HI	381
B-4	Cl	CF <sub>3</sub>	H	H	Me	H	H	H	H	359
B-5	Cl	CF <sub>3</sub>	H	H	F	Cl	H	H	H	398

The following examples of compound preparation are mentioned with the aim of illustrating the invention, but should in no way be considered to limit the said invention.


5 **Preparation of methyl [3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)acetate:**



Procedure:

Under argon, 116 g of sodium hydride, 60% in dispersion in oil (2.91 mol, 1.8 eq.), are suspended in 3 L of DMF. The suspension is cooled in a bath of ice cold water. 160 g (1.616 mol, 1.0 eq.) of methyl cyanoacetate in solution in 200 mL of DMF are added dropwise, with stirring. Once all gas has been given off, 350 g (1.616 mol, 1.0 eq) of 2,3-dichloro-5-(trifluoromethyl)pyridine are added with stirring. The mixture is stirred overnight at ambient temperature. 50 mL of methanol are added. The reaction medium is poured into 5 L of water. The pH is adjusted to 3-4 with concentrated hydrochloric acid. The yellow precipitate of methyl [3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)acetate which forms is filtered off and washed with water and with pentane.

20 **Preparation of [3-chloro-5-(trifluoromethyl)-2-pyridinyl]acetonitrile:**




Procedure:

314 g (1.13 mol, 1 eq.) of methyl [3-chloro-5-(trifluoromethyl)-2-pyridinyl]-(cyano)acetate and 22 g (0.38 mol, 0.33 eq.) of sodium chloride are dissolved in a solution of 44 mL of water and 1.1 L of dimethyl sulphoxide. The reaction medium is stirred and heated at 160°C. Once all gas has been given off, the medium is cooled to ambient temperature. 1 L of water and 0.5 L of dichloromethane are added. After separation, the aqueous phase is extracted twice with 0.5 L of dichloromethane. The organic phase is washed twice with 0.5 L of water and dried over magnesium



5 sulphate. After concentration, the crude product is diluted in 100 mL of dichloromethane and eluted with an ethyl acetate/heptane mixture (20/80) on a bed of silica. The filtrate is concentrated so as to produce [3-chloro-5-(trifluoromethyl)-2-pyridinyl]acetonitrile.


**Preparation of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl] ethanamine acetate:**



Procedure:

- 10 113 g of [3-chloro-5-(trifluoromethyl)-2-pyridinyl]acetonitrile (0.51 mol, 1 eq.) are diluted in 2.5 L of acetic acid. 30 g of palladium (5% on charcoal) are added. The reaction medium is stirred at ambient temperature under a hydrogen pressure of 5 bar. The progress of the reaction is followed by TLC; when the [3-chloro-5-(trifluoromethyl)-2-pyridinyl]acetonitrile has been completely used up, the medium is filtered over a bed of celite, and then concentrated to dryness so as to produce the 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl] ethanamine acetate.

**Preparation of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-benzamides:**



Procedure:

- 20 0.100 g of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl] ethanamine acetate (0.00037 mol, 1.0 eq.) are diluted in 3 mL of dichloromethane. 0.500 g of poly(4-vinylpyridine) are added. The mixture is stirred at ambient temperature for half an hour.

25 1.2 equivalents of the desired acyl chloride are added. The reaction medium is stirred at ambient temperature overnight, filtered, and concentrated to dryness. The crude product is then purified by reverse-phase HPLC. The corresponding amide is obtained.

The following biological activities were tested in order to establish a comparison between the activity of the compounds of general formula (I) according to the present invention and a compound described in patent application WO 01/11965, on a certain number of fungal diseases.

In vivo test of activity on *Alternaria brassicae* (crucifer black spot)

An aqueous solution of the active material to be tested is prepared at a concentration of 2 g/l by grinding in solution 1 or solution 2:

Solution 1

- Water
- Tween 80 diluted to 10% in water: 5 µl/mg of active material
- Clay in a sufficient amount for active material + clay to equal 100 mg;

Solution 2

Concentrated solution containing surfactants and adjuvants conventionally used: 200 µl/mg of active material.

The aqueous solution is diluted with water so as to obtain the desired concentration.

Radishes of the Pernot variety are sown in a 50/50 pozzolan/peat substrate and maintained at 18-22°C. The treatment is carried out by spraying the aqueous suspension. The untreated control plants are sprayed with water. 24 hours after treatment, the plants are inoculated by spraying a solution of *Alternaria brassicae* spores (40 000 spores/ml) originating from a 12-day culture.

The radish plants are then maintained at 18-20°C in a humid atmosphere. The effectiveness of the products is evaluated relative to the control plants after seven to eight days of incubation.

In vivo test of activity on *Botrytis cinerea* on cucumber

An aqueous solution of the active material to be tested is prepared at a concentration of 2 g/l by grinding in solution 1 or solution 2:

Solution 1

- Water
- Tween 80 diluted to 10% in water: 5 µl/mg of active material
- Clay in a sufficient amount for active material + clay to equal 100 mg;

Solution 2

Concentrated solution containing surfactants and adjuvants conventionally used: 200 µl/mg of active material.

5        The aqueous solution is diluted with water so as to obtain the desired concentration.

Cucumbers of the Marketer variety are sown in a 50/50 pozzolan/peat substrate and maintained at 18-22°C. The treatment is carried out by spraying the aqueous suspension. The untreated control plants are sprayed with water.

10       24 hours after treatment, the plants are inoculated by spraying a solution of *Botrytis cinerea* spores (150 000 spores/ml) originating from a 15-day culture.

The cucumber plants are then maintained at 11-15°C in a humid atmosphere. The effectiveness of the products is evaluated relative to the control plants after seven to eight days of incubation.

15

In vivo test of activity on *Pyrenophora teres* (barley net blotch)

An aqueous solution of the active material to be tested is prepared at a concentration of 2 g/l by grinding in solution 1 or solution 2:

Solution 1

- 20       -        Water  
         -        Tween 80 diluted to 10% in water: 5 µl/mg of active material  
         -        Clay in a sufficient amount for active material + clay to equal 100 mg;

Solution 2

25       Concentrated solution containing surfactants and adjuvants conventionally used: 200 µl/mg of active material.

The aqueous solution is diluted with water so as to obtain the desired concentration.

30       Barley plants of the Express variety are sown in a 50/50 pozzolan/peat substrate and maintained at 12°C. The treatment is carried out at the 1-leaf stage (10 cm) by spraying the aqueous suspension. The untreated control plants are sprayed with water.

24 hours after treatment, the plants are inoculated by spraying a solution of *Pyrenophora teres* spores (10 000 spores/ml) originating from a 10-day culture.

35       The barley plants are then maintained at 18°C in a humid atmosphere. The effectiveness of the products is evaluated relative to the control plants after eight to

fifteen days of incubation.

In vivo test of activity on *Septoria tritici* (wheat septoria disease)

5 An aqueous solution of the active material to be tested is prepared at a concentration of 2 g/l by grinding in solution 1 or solution 2:

Solution 1

- Water
- Tween 80 diluted to 10% in water: 5 µl/mg of active material
- Clay in a sufficient amount for active material + clay to equal 100 mg;

10 Solution 2

Concentrated solution containing surfactants and adjuvants conventionally used: 200 µl/mg of active material.

15 The aqueous solution is diluted with water so as to obtain the desired concentration.

Wheat plants of the Scipion variety are sown in a 50/50 pozzolan/peat substrate and maintained at 12°C. The treatment is carried out at the 1-leaf stage (10 cm) by spraying the aqueous suspension. The untreated control plants are sprayed with water.

20 24 hours after treatment, the plants are inoculated by spraying a solution of *Septoria tritici* spores (500 000 spores/ml) from a 7-day culture.

The wheat plantlets are then maintained at 18-20°C in a humid atmosphere for 72 hours, and then at 90% relative humidity. The effectiveness of the products is evaluated relative to the control plants, 21 to 28 days after the contamination.

25

The effectiveness of the molecules is estimated, at 500 g/ha, 250 g/ha or 330 ppm, by the percentage of control relative to untreated plants. Under these conditions, good effectiveness is defined as more than 80% effectiveness. Average effectiveness is defined as an effectiveness between 50 and 80%. Poor effectiveness is defined as an effectiveness between 10 and 50% and zero effectiveness is defined as less than 10% effectiveness.

30

At a concentration of 500 g/ha, the following compounds showed good to average effectiveness against the fungal pathogens:

35 *Alternaria brassicae* : A-2, A-4, A-6, A-7, A-9, A-13, A-14, A-20, A-25.

*Botrytis cinerea* : A-2, A-7, A-9, A-20, A-25.

*Pyrenophora teres* : A-2, A-4, A-5, A-6, A-7, A-9, A-20, A-25, A-27.

*Septoria tritici* : A-2, A-4, A-5, A-6, A-7, A-16, A-18, A-20, A-21, A-22, A-23, A-24, A-25.

5           At a concentration of 250 g/ha, the following compounds showed good to average effectiveness against the fungal pathogens:

*Alternaria brassicae* : A-20, A-28, A-29, A-41, A-45, A-46, A-73, A-173.

*Botrytis cinerea* : A-20, A-45, A-46, A-73.

*Pyrenophora teres* : A-20, A-45, A-46, A-73.

10

          At a concentration of 330 ppm, the following compounds showed good to average effectiveness against the fungal pathogens:

*Alternaria brassicae* : A-20, A-44, A-45, A-46, A-47, A-48, A-49, A-52, A-60, A-61, A-62, A-71, A-72, A-73, A-74, A-75, A-76, A-77, A-79, A-80, A-83, A-84, A-85, A-86, A-87, A-89, A-91, A-92, A-96, A-98, A-99, A-100, A-107, A-110, A-112, A-113, A-117, A-122, A-123, A-124, A-125, A-127, A-128, A-133, A-134, A-135, A-136, A-137, A-138, A-139, A-140, A-141, A-142, A-143, A-144, A-146, A-147, A-148, A-150, A-151, A-152, A-156, A-157, A-158, A-159, A-162, A-165, A-166, A-167, A-168, A-169, A-170, A-171, A-173, A-174, A-175, A-176, A-177, A-178, A-179, A-180, A-181, A-182, A-183, A-184, A-185, A-186, A-187, A-188, A-189, A-194, B-1.

20

*Botrytis cinerea* : A-20, A-45, A-46, A-73, A-170, A-172, A-173, A-174, A-175, A-187.

*Pyrenophora teres* : A-20, A-44, A-45, A-46, A-61, A-73, A-83, A-87, A-89, A-96, A-117, A-125, A-133, A-134, A-140, A-167, A-173, A-174, A-187, B-1.

25

          Under these conditions, the N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-ethyl}-2,6-dichlorobenzamide showed respectively poor and zero effectiveness on *Alternaria Brassicae* at 330ppm and 250 g/ha; and zero effectiveness on *Botrytis cinerea* at 250 g/ha and 330ppm.

30

          Under these conditions, the N-{1-methylcarbamoyl-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-4-phenylbenzamide disclosed by Patent Application WO 01/11965 (see compound 316 in Table D) showed poor effectiveness on *Alternaria Brassicae* and *Septoria tritici*, and zero effectiveness on *Botrytis cinerea* at 250 g/ha; the N-{1-ethylcarbamoyl-2-[3-chloro-


35



5-(trifluoromethyl)-2-pyridinyl]ethyl}-3-nitrobenzamide also disclosed by Patent Application WO 01/11965 (see compound 307 in Table D) showed zero effectiveness on *Alternaria Brassicae* and *Botrytis cinerea* at 250 g/ha; the N-{1-ethylcarbamoyl-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-benzamide and  
5 the N-{1-methylcarbamoyl-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-benzamide also disclosed by Patent Application WO 01/11965 (see compounds 304 and 314 in Table D) showed poor effectiveness on *Septoria tritici* and zero effectiveness on *Botrytis cinerea* at 250 g/ha; and the N-{1-ethylcarbamoyl-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-4-chlorobenzamide, the N-{1-ethylcarbamoyl-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-  
10 bromobenzamide and the N-{1-methylcarbamoyl-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-4-methoxybenzamide also disclosed by Patent Application WO 01/11965 (see compounds 306, 310 and 315 in Table D) showed zero effectiveness on *Botrytis cinerea* at 250 g/ha.

**CLAIMS**

1. Compound of general formula (I):




in which

- 5 - p is an integer equal to 1, 2, 3 or 4;  
 - q is an integer equal to 1, 2, 3, 4 or 5;  
 - each substituent X is chosen, independently of the others, as being halogen, alkyl or haloalkyl, at least one of the substituents being a haloalkyl;  
 - each substituent Y is chosen, independently of the others, as being halogen, alkyl,  
 10 alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl, alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or benzylsulphonyl;  
 as to the N-oxides of 2-pyridine thereof;  
 15 with the exception of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2,6-dichlorobenzamide.


2. Compound according to Claim 1, characterised in that p is equal to 2.

- 20 3. Compound according to Claim 2, characterised in that the substituents X are positioned as follows:




4. Compound according to any one of Claims 1 to 3, characterised in that q is chosen equal to 1 or 2, the substituent(s) Y being positioned in the ortho position of  
 25 the benzene ring.

5. Compound according to Claim 4, characterised in that it corresponds to general formula (I'):



6. Compound according to Claim 5, characterised in that X<sup>1</sup> is halogen and X<sup>2</sup> is haloalkyl.

7. Compound according to Claim 4, characterised in that it corresponds to general formula (I''):



8. Compound according to Claim 7, characterised in that it has the following characteristics, taken individually or in combination:

- X<sup>1</sup> is chosen as being halogen and X<sup>2</sup> is chosen as being haloalkyl;
- Y<sup>1</sup> is chosen as being halogen or haloalkyl.

9. Compound according to Claim 8, characterised in that haloalkyl group is trifluoromethyl.


10. Compound according to Claim 9, characterised in that the compound of formula (I'') is:

- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide;
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide; or
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-bromobenzamide.

11. Process for the preparation of a compound according to any one of Claims 1 to 10, characterised in that it comprises :

- a first step consisting in reacting, in the presence of a base in aprotic polar solvent, a compound of general formula (Ia) in order to substitute it selectively in the 2-position:


\* either with a group of the alkyl cyanoacetate type (NC-CH<sub>2</sub>-CO<sub>2</sub>Alk) to produce a compound of general formula (Ib) according to the following reaction scheme:




where: - X is as defined where Y is as defined in any one of Claims 1 to 9;

- Alk represents an alkyl radical;  
 - Q is a nucleofugal radical;

the compound of general formula (Ib) thus obtained then undergoing dealkyloxy-carbonylation in the presence of an alkali metal halide, such as Li-halogen, K-halogen or Na-halogen, at the reflux of a water/dimethyl sulphoxide mixture, according to the Krapcho reaction to produce the compound of general formula (Ic) according to the following reaction scheme:




\* or with acetonitrile to directly produce the compound of general formula (Ic) according to the following reaction scheme:




- a second step consisting in the reduction of the compound of general formula (Ic) to

pyridylethanamine of general formula (Id) (or its corresponding ammonium salt depending on whether or not the medium is acid) under hydrogen pressure in the presence of a metal catalyst in a protic solvent according to the following reaction scheme:



- 5 - a third step consisting in converting the compound of general formula (Id) to a compound of general formula (I) by reaction with a benzoyl halide in the presence of a base according to the following reaction scheme:



where Y is as defined in any one of Claims 1 to 9.

- 10 **12.** Process according to Claim 11, characterised in that the nucleofugal radical Q is a halogen or trifluoromethanesulphonate.
- 13.** Fungicidal composition comprising an effective amount of a compound according to any of the claims 1 to 10 and an agriculturally acceptable support.
- 15 **14.** Fungicidal composition according to claim 13 further comprising a surfactant.
- 15.** Fungicidal composition according to either of claims 13 and 14, comprising from 0.05% to 99% by weight of active material.
- 20 **16.** Method for preventively or curatively combating the phytopathogenic fungi of crops, characterised in that an effective and non-phytotoxic amount of a composition according to any of the claims 13 to 15 is applied to the plant seeds or to the plant leaves and/or to the fruits of the plants or to the soil in which the plants are



growing or in which it is desired to grow them.